



Cucurbitae Pro Drop®



Solution de piégeage basée sur un attractant alimentaire contre *Bactrocera cucurbitae* et *Dacus demmerezii*

Cette solution combine un Smart trap® et un diffuseur d'attractif alimentaire pour capturer les mâles et les femelles adultes des mouches des cucurbitacées.

Avantages



Diffusion longue durée



Formulé avec des ingrédients biosourcés



Utilisable en agriculture biologique



Facile à appliquer



Outil d'aide à la décision



Spécifique au ravageur ciblé



Stockage à température ambiante (0-30°C)

Caractéristiques de Cucurbitae Pro Drop®

Type de produit	Diffuseur d'attractant alimentaire
Conditionnement	Tube Eppendorf
Usage	Détection / Monitoring et piégeage de masse
Substance active	Cuelure
Cible	les mouches des cucurbitacées (<i>Bactrocera cucurbitae</i> et <i>Dacus demmerezii</i>)
Volume de formulation	1,45 g
Durée indicative de diffusion*	4 à 8 semaines
Rayon estimé de diffusion	Mouches attirées sur un rayon d'environ de 10 m

* pour une température moyenne de 30 °C et sans vent fort.

Utilisation

- 🕒 **Période de détection** : sur toute l'année. Renouveler le diffuseur selon la durée indicative de diffusion. Effectuer des relevés de manière hebdomadaire pour suivre la pression du ravageur.
- 📍 **Positionnement du piège** : Suspendre dans un arbre à 1,5 m du sol.
- 📊 **Densité recommandée pour le monitoring** : 2- 5 pièges/ha
- 📊 **Densité recommandée pour le piégeage en masse** : 50 pièges/ha. 14 mètres entre chaque piège
- 📖 Consultez votre agronome de référence pour plus de détails.



Pièges associés



Montage de piège

Exemple d'utilisation de **Cucurbitae Pro Drop®** + Smart Trap® (format à l'envers)



Percez des trous dans le bol pour y faire passer la ficelle. Porter des gants. Ouvrir le tube d'attractif. Ne pas toucher le produit avec les mains



Placer le tube (ouverture vers le haut) dans la cage. Fixer la cage dans le bol (supports fournis)



Ajouter de l'eau ou une solution de savon inodore dans la partie inversée (partie verte). Tordre ensemble le bol et l'entonnoir. Suspendre le piège à l'envers. Les mouches mâles et femelles sont attirées, entrent dans le piège et se noient dans l'eau.

Consultez notre catalogue complet





Trapping solution based on food attractant against *Bactrocera cucurbitae* and *Dacus demmerezi*

This solution combines a Smart trap® and a food attractant diffuser to capture the males and females adults of of cucurbit flies.

Benefits



Long diffusion time



Formulated with bio-based ingredients



Usable in organic farming



Easy to apply



Decision support tool



Species-specific targeting



Storage at room temperature (0-30°C)

Characteristics of Cucurbitae Pro Drop®

Product type	Food attractant dispenser
Packaging	Eppendorf tubes
Use	Detection / Monitoring & Mass trapping
Active substance	Cuelure
Target	the Cucurbit flies (<i>Bactrocera cucurbitae</i> and <i>Dacus demmerezi</i>)
Volume of formulation	1,45 g
Indicative diffusion time*	4 to 8 weeks
Estimated diffusion radius	Flies attracted on a radius of approximatively 10 m

* at an average temperature of 30 °C and without strong wind.

Use

- ☞ **Detection period:** throughout the season. Renew the dispenser according to the indicative duration of the broadcast. Take weekly readings to monitor pest pressure and empty the trap if necessary.
- ☞ **Trap location:** Hung trap in a tree at 1.5 m from the ground.
- ☞ **Recommended density for monitoring:** 2-5 traps/ha
- ☞ **Recommended density for mass trapping:** 50 traps/ha. 14m between each traps.
- ☞ Consult your reference agronomist for more details



Associated traps



Trap setup

Example of use of an **Cucurbitae Pro Drop®** + Smart Trap® (Upside down format)



Drill holes in the bowl to pass the string through. Wear gloves. Open the tube of attractant. Do not touch the product with your hands



Place the tube (opening upwards) in the cage. Attach the cage in the bowl (supports provided)



Add water or an odorless soap solution in the upside-down (green part). Twist together the bowl and the funnel. Hang the trap in the upside-down position. Male and female flies are attracted, enter the trap and drown in the water.

