



Solution de piégeage à base de phéromone contre *Cydia pomonella*

Cette solution combine un Delta trap et un diffuseur à base de phéromone sexuelle pour capturer les mâles adultes du carpocapse des pommes et des poires.

Avantages



Diffusion longue durée



Formulé avec des ingrédients biosourcés



Utilisable en agriculture biologique



Facile à appliquer



Outil d'aide à la décision



Spécifique au ravageur ciblé



Stockage à température ambiante (0-30°C)

Caractéristiques de Cydia Pro Caps®

Type de produit	Diffuseur de phéromone
Packaging	Seringue en polypropylène
Utilisation	Détection / Monitoring
Substance active	Codlemone
Cible	Carpocapse des pommes et des poires (<i>Cydia pomonella</i>)
Poids net de la formulation	1,08 g
Temps indicatif de diffusion*	3 mois
Rayon de diffusion estimé	Papillons attirés sur un rayon de 5 à 10 m

*pour une température moyenne de 30°C et sans vent fort.

Utilisation

- 🕒 **Période de détection** : de mi-avril à septembre. Renouvelez le diffuseur en fonction de la durée indicative de diffusion. Effectuez des relevés de manière hebdomadaire pour suivre la pression du ravageur.
- 📍 **Positionnement des pièges** : suspendre dans la partie supérieure du feuillage.
- 📊 **Densité recommandée pour du monitoring** : 1 piège/ha.
- 📖 Consultez votre agronome référent pour plus de détails.



Pièges associés




Montage du piège

Exemple d'utilisation de la seringue **Cydia Pro Caps®** + Delta trap




Montez le piège Delta



Retirer le film protecteur de la plaque engluée.



Placer la coupelle verte au milieu la plaque engluée. Vider le contenu de la seringue par simple pression sur le piston (inutile de découper l'embout) dans la coupelle.



Insérer la plaque dans le piège avant de le refermer.





Trapping solution based on pheromone against *Cydia pomonella*

This solution combines a Delta trap and a sexual pheromone attractant dispenser to capture the males' adults of the codling moth.

Benefits



Long diffusion time



Formulated with bio-based ingredients



Usable in organic farming



Easy to apply



Decision support tool



Species-specific targeting



Storage at room temperature (0-30°C)

Characteristics of Cydia Pro Caps®

Product type	Pheromone-attractant dispenser
Packaging	Polypropylene syringe
Use	Detection / Monitoring
Active substance	Codlemone
Target	Codling moth (<i>Cydia pomonella</i>)
Net weight of formulation	1,08 g
Indicative diffusion time*	3 months
Estimated diffusion radius	Moths attracted on a radius from 5 to 10 m

*at an average temperature of 30°C and without strong wind.

Use

- 🕒 **Detection period:** from mid-April to September (adapt and renew the dispenser according to recommended diffusion time).
- 📍 **Trap location:** Hung in the upper part of the foliage.
- 📊 **Recommended density for monitoring:** 1 to 2 trap/ha.
- 📖 Consult your reference agronomist for more details



Associated traps




Trap setup

Example of use of **Cydia Pro Caps® syringe** + Delta trap




Mount the Delta Trap



Remove the protective film from the glue plate



Place the green cup in the middle of the plate and empty the syringe contents into the cup by simply pressing the plunger (no need to cut the tip).



Insert the plate into the trap before closing it

