



Solution de monitoring à base de phéromones contre *Grapholita funebrana*

Cette solution combine un piège Delta et un diffuseur à base de phéromone pour capturer les mâles adultes du carpocapse de prune.

Avantages



Diffusion longue durée



Formulé avec des ingrédients biosourcés



Utilisable en agriculture biologique



Facile à appliquer



Outil d'aide à la décision



Spécifique au ravageur ciblé



Stockage à température ambiante (0-30°C)

Caractéristiques de Funebrana Pro Caps®

Type de produit	Diffuseur de phéromones
Conditionnement	seringue en propylène
Usage	Détection / Monitoring
Substance active	Z-8 dodecen-1-yl acetate ; E-8 dodecen-1-yl acetate
Cible	Carpocapse des prunes (<i>Grapholita funebrana</i>)
Volume de formulation	0,5 mL
Durée indicative de diffusion*	3 mois
Rayon estimé de diffusion	Papillons attirés sur un rayon d'environ de 5 à 10 m

* pour une température moyenne de 30 °C et sans vent fort.

Utilisation

- ⊖ **Période de détection** : d'Avril à octobre (adapter et renouveler le diffuseur de phéromones en fonction de la durée indicative de diffusion).
- ⊖ **Positionnement du piège** : suspendu à environ 1,5 m au-dessus de la culture.
- ⊖ **Densité recommandée pour le monitoring** : 3 à 5 pièges/ha (Frasin et al, 2010).
- ⊖ Consultez votre agronome de référence pour plus de détails.



Pièges associés



Delta Trap®
Recommandé



Smart Trap®
Alternative

Montage de piège

Exemple d'utilisation de Funebrana Pro Caps® + Delta trap




Montez le piège Delta



Retirer le film protecteur de la plaque engluée.



Placer la coupelle verte au milieu la plaque engluée. Vider le contenu de la seringue par simple pression sur le piston (inutile de découper l'embout) dans la coupelle.



Insérer la plaque dans le piège avant de le refermer.





Monitoring solution based on pheromone against *Grapholita funebrana*

This solution combines a Delta trap and a pheromone attractant dispenser to capture Plum fruit moth adults.

Benefits



Long diffusion time



Formulated with bio-based ingredients



Usable in organic farming



Easy to apply



Decision support tool



Species-specific targeting



Storage at room temperature (0-30°C)

Characteristics of Funebrana Pro Caps®

Product type	Food attractant dispenser
Packaging	Polypropylene syringe
Use	Detection / Monitoring
Active substance	Z-8 dodecen-1-yl acetate ; E-8 dodecen-1-yl acetate
Target	Plum Fruit moth (<i>Grapholita funebrana</i>)
Volume of formulation	0,5 mL
Indicative diffusion time*	3 months
Estimated diffusion radius	Moths attracted on a radius of approximately of 5-10 m

* at an average temperature of 30 °C and without strong wind.

Use

- 🕒 **Detection period:** from April to October (adapt and renew the pheromone dispenser according to the recommended diffusion time).
- 📍 **Trap location:** hung on the tree's canopy, 1,5m from the ground.
- 👤 **Recommended density for monitoring:** 3-5 traps/ha (Frasin et al., 2010).
- 📖 Consult your reference agronomist for more details



Associated traps




Trap setup

Example of use of an Funebrana Pro Caps® + Delta trap





Mount the Delta Trap



Remove the protective film from the glue plate



Place the green cup in the middle of the plate and empty the syringe contents into the cup by simply pressing the plunger (no need to cut the tip).



Insert the plate into the trap before closing it

Check out our full catalog

