



Solution de piégeage à base de phéromone contre *Plutella xylostella*

Cette solution combine un Delta trap et un diffuseur à base de phéromone sexuelle pour capturer les mâles adultes de la teigne des crucifères.

Avantages



Diffusion longue durée



Formulé avec des ingrédients biosourcés



Utilisable en agriculture biologique



Facile à appliquer



Outil d'aide à la décision



Spécifique au ravageur ciblé



Stockage à température ambiante (0-30°C)

Caractéristiques de Plutella Pro Caps®

Type de produit	Diffuseur de phéromone
Packaging	Seringue en polypropylène
Utilisation	Détection / Monitoring
Substance active	Z11-hexadecenyl acetate; Z11-hexadecenol; Z11-hexadecenal
Cible	Teigne des crucifères (<i>Plutella xylostella</i>)
Poids net de formulation	0,45 g
Temps indicatif de diffusion*	2 mois
Rayon de diffusion estimé	Papillons attirés sur un rayon de 5 à 10 m

*pour une température moyenne de 30°C et sans vent fort.

Utilisation

- 🕒 **Période de détection** : d'avril à octobre. Renouvelez le diffuseur en fonction de la durée indicative de diffusion. Effectuez des relevés de manière hebdomadaire pour suivre la pression du ravageur.
- 📍 **Positionnement des pièges** : suspendre au-dessus du sol.
- 📊 **Densité recommandée pour du monitoring** : 4-6 pièges/ha ou 2 pièges/500 m² sous serre.
- 📖 Consultez votre agronome référent pour plus de détails.




Pièges associés



Montage du piège

Exemple d'utilisation de la **seringue Plutella Pro Caps®** + Smart Trap®





Trapping solution based on pheromone against *Plutella xylostella*

This solution combines a Delta trap and a sexual pheromone attractant dispenser to capture the males' adults of the Diamondback moth.

Benefits



Long diffusion time



Formulated with bio-based ingredients



Usable in organic farming



Easy to apply



Decision support tool



Species-specific targeting



Storage at room temperature (0-30°C)

Characteristics of Plutella Pro Caps®

Product type	Pheromone-attractant dispenser
Packaging	Polypropylene syringe
Use	Detection / Monitoring
Active substance	Z11-hexadecenyl acetate; Z11-hexadecenol; Z11-hexadecenal
Target	Diamond black moth (<i>Plutella xylostella</i>)
Net weight of formulation	0,45 g
Indicative diffusion time*	2 months
Estimated diffusion radius	Moths attracted on a radius from 5 to 10 m


*at an average temperature of 30°C and without strong wind.

Use

- 🕒 **Detection period:** from April to October. Renew the dispenser according to the indicative diffusion time. Conduct weekly the traps to monitor pest pressure.
- 📍 **Trap location:** hung above the crop.
- 📊 **Recommended density for monitoring:** 4-6 traps/ha or 2 traps/500 m² in greenhouses.
- 📊 **Recommended density for mass trapping:** 60 traps/ha, spaced 13 m apart.
- 📖 Consult your reference agronomist for more details.



Associated traps



Trap setup

Example of use of **Plutella Pro Caps® syringe** + Smart Trap®



Drop a drop at the bottom of the bowl by simply pressing the plunger (there is no need to cut the tip).



Empty the rest of the syringe contents into the cage and close it.



Make sure that the cage is placed in the hole provided for this purpose in the roof of the trap.

Do not add water to the trap bowl!!

Assemble the two parts of the trap by twisting them together



Male butterflies are attracted to the trap



Check out our full catalog

